

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

A Prospective International Aspergillus terreus Survey: An EFISG, ISHAM and ECMM Joint Study

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1638082> since 2017-10-09T18:02:32Z

Published version:

DOI:doi.org/10.1016/j.cmi.2017.04.012

Terms of use:

Open Access


Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Accepted Manuscript

A Prospective International *Aspergillus terreus* Survey: An EFISG, ISHAM and ECMM Joint Study

Brigitte Risslegger, Tamara Zoran, Michaela Lackner, Maria Aigner, Ferran Sánchez-Reus, Antonio Rezusta, Anuradha Chowdhary, Saad Jaber Taj-Aldeen, Maiken C. Arendrup, Salvatore Oliveri, Dimitrios P. Kontoyiannis, Ana Alastruey-Izquierdo, Katrien Lagrou, Giuliana Lo Cascio, Jacques F. Meis, Walter Buzina, Claudio Farina, Miranda Drogari-Apiranthitou, Anna Grancini, Anna Maria Tortorano, Birgit Willinger, Axel Hamprecht, Elizabeth Johnson, Lena Klingspor, Valentina Arsic-Arsenijevic, Oliver A. Cornely, Joseph Meletiadis, Wolfgang Prammer, Vivian Tullio, Jörg-Janne Vehreschild, Laura Trovato, Russell E. Lewis, Esther Segal, Peter-Michael Rath, Petr Hamal, Manuel Rodriguez-Iglesias, Emmanuel Roilides, Sevtap Arikan-Akdogli, Arunaloke Chakrabarti, Arnaldo L. Colombo, Mariana S. Fernández, M. Teresa Martin-Gomez, Hamid Badali, Georgios Petrikos, Nikolai Klimko, Sebastian M. Heimann, Jos Houbbraken, Omrum Uzun, Michael Edlinger, Sonia de la Fuente, Cornelia Lass-Flörl



PII: S1198-743X(17)30220-3

DOI: [10.1016/j.cmi.2017.04.012](https://doi.org/10.1016/j.cmi.2017.04.012)

Reference: CMI 924

To appear in: *Clinical Microbiology and Infection*

Received Date: 24 February 2017

Revised Date: 7 April 2017

Accepted Date: 9 April 2017

Please cite this article as: Risslegger B, Zoran T, Lackner M, Aigner M, Sánchez-Reus F, Rezusta A, Chowdhary A, Taj-Aldeen SJ, Arendrup MC, Oliveri S, Kontoyiannis DP, Alastruey-Izquierdo A, Lagrou K, Lo Cascio G, Meis JF, Buzina W, Farina C, Drogari-Apiranthitou M, Grancini A, Tortorano AM, Willinger B, Hamprecht A, Johnson E, Klingspor L, Arsic-Arsenijevic V, Cornely OA, Meletiadis J, Prammer W, Tullio V, Vehreschild J-J, Trovato L, Lewis RE, Segal E, Rath P-M, Hamal P, Rodriguez-Iglesias M, Roilides E, Arikan-Akdogli S, Chakrabarti A, Colombo AL, Fernández MS, Martin-Gomez MT, Badali H, Petrikos G, Klimko N, Heimann SM, Houbbraken J, Uzun O, Edlinger M, Fuente Sdl, Lass-Flörl C, A Prospective International *Aspergillus terreus* Survey: An EFISG, ISHAM and ECMM Joint Study, *Clinical Microbiology and Infection* (2017), doi: 10.1016/j.cmi.2017.04.012.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

RESEARCH NOTE

A Prospective International *Aspergillus terreus* Survey:
An EFISG, ISHAM and ECMM Joint Study

Brigitte Risslegger^{1*}, Tamara Zoran^{1*}, Michaela Lackner¹,
Maria Aigner¹, Ferran Sánchez-Reus², Antonio Rezusta³,
Anuradha Chowdhary⁴, Saad Jaber Taj-Aldeen⁵, Maiken C.
Arendrup⁶, Salvatore Oliveri⁷, Dimitrios P. Kontoyiannis⁸, Ana
Alastruey-Izquierdo⁹, Katrien Lagrou¹⁰, Giuliana Lo Cascio¹¹,
Jacques F. Meis¹², Walter Buzina¹³, Claudio Farina¹⁴, Miranda
Drogari-Apiranthitou¹⁵, Anna Grancini¹⁶, Anna Maria
Tortorano¹⁷, Birgit Willinger¹⁸, Axel Hamprecht¹⁹, Elizabeth
Johnson²⁰, Lena Klingspor²¹, Valentina Arsic-Arsenijevic²²,
Oliver A. Cornely²³, Joseph Meletiadis²⁴, Wolfgang Prammer²⁵,
Vivian Tullio²⁶, Jörg-Janne Vehreschild²⁷, Laura Trovato²⁸,
Russell E. Lewis²⁹, Esther Segal³⁰, Peter-Michael Rath³¹, Petr
Hamal³², Manuel Rodriguez-Iglesias³³, Emmanuel Roilides³⁴,
Sevtap Arikan-Akdagli³⁵, Arunaloke Chakrabarti³⁶, Arnaldo L.
Colombo³⁷, Mariana S. Fernández³⁸, M. Teresa Martin-
Gomez³⁹, Hamid Badali⁴⁰, Georgios Petrikos⁴¹, Nikolai
Klimko⁴², Sebastian M. Heimann⁴³, Jos Houbraken⁴⁴, Omrum

24 Uzun⁴⁵, Michael Edlinger⁴⁶, Sonia de la Fuente⁴⁷, Cornelia Lass-

25 Flörl¹

26

27 * contributed equally

28

29 Author affiliations:

30 ¹Division of Hygiene and Medical Microbiology, Medical

31 University of Innsbruck, Innsbruck/Austria, ²Servei de

32 Microbiologia, Hospital de la Santa Creu I Sant Pau,

33 Barcelona/Spain, ³Microbiologia, Hospital Universitario Miguel

34 Servet, IIS Aragon, Universidad de Zaragoza, Zaragoza/Spain,

35 ⁴Department of Medical Mycology, Vallabhbhai Patel Chest

36 Institute, University of Delhi, Delhi/India, ⁵Microbiology

37 Division, Department of Laboratory Medicine and Pathology,

38 Hamad Medical Corporation, Doha/Qatar, ⁶Statens Serum

39 Institute, Unit of Mycology, Copenhagen/Denmark &

40 Department of Clinical Microbiology, Copenhagen University,

41 Rigshospitalet, Copenhagen/Denmark, ⁷Department of

42 Biomedical and Biotechnological Sciences, University of

43 Catania, Catania/Italy, ⁸The University of Texas MD Anderson

44 Cancer Center, Houston, Texas/USA, ⁹National Centre for

45 Microbiology, Instituto de Salud Carlos III, Madrid/Spain,

46 ¹⁰Department of Microbiology and Immunology, KU Leuven,

47 Leuven/Belgium, ¹¹Unità Operativa Complessa di Microbiologia

48 e virologia, Dipartimento di Patologia e diagnostica, Azienda
 49 Ospedaliera Universitaria Integrata, Verona/Italy,
 50 ¹²Department of Medical Microbiology and Infectious
 51 Diseases, Canisius Wilhelmina Hospital, Nijmegen/The
 52 Netherlands, ¹³Institute of Hygiene, Microbiology and
 53 Environmental Medicine, Medical University of Graz,
 54 Graz/Austria, ¹⁴Microbiology Institute, ASST Papa Giovanni
 55 XXIII, Bergamo/Italy, ¹⁵Infectious Diseases Research
 56 Laboratory/⁴th Department of Internal Medicine, ATTIKON
 57 University Hospital, National and Kapodistrian University of
 58 Athens, Athens/Greece, ¹⁶Laboratorio Centrale di Analisi
 59 Chimico Cliniche e Microbiologia, IRCCS Foundation, Cà
 60 Granda Ospedale Maggiore Policlinico, Milan/Italy,
 61 ¹⁷Department of Biomedical Sciences for Health, Università
 62 degli Studi di Milano, Milan/Italy, ¹⁸Department of Laboratory
 63 Medicine, Division of Clinical Microbiology, Medical University
 64 of Vienna, Vienna/Austria, ¹⁹Institute for Medical
 65 Microbiology, Immunology and Hygiene, University of
 66 Cologne, Cologne/Germany, ²⁰Mycology Reference Laboratory,
 67 Public Health England, Bristol/United Kingdom, ²¹Karolinska
 68 Institutet, Department of Laboratory Medicine, F 68,
 69 Karolinska University Hospital, Huddinge, Stockholm/Sweden,
 70 ²²National Reference Medical Mycology Laboratory, Institute
 71 of Microbiology and Immunology, Faculty of Medicine,

72 University of Belgrade, Belgrade/Serbia, ²³Cologne Excellence
 73 Cluster on Cellular Stress Responses in Aging-Associated
 74 Diseases (CECAD), Department I of Internal Medicine, Clinical
 75 Trials Centre Cologne (ZKS Köln), Center for Integrated
 76 Oncology (CIO Köln-Bonn), German Centre for Infection
 77 Research (DZIF), University of Cologne, Cologne/Germany,
 78 ²⁴Clinical Microbiology Laboratory, National Kapodistrian
 79 University of Athens, ATTIKON University Hospital Athens,
 80 Athens/Greece, ²⁵Department of Hygiene and Medical
 81 Microbiology, Klinikum Wels-Grieskirchen, Wels/Austria,
 82 ²⁶Department of Public Health and Pediatrics, Microbiology
 83 Division, Turin/Italy, ²⁷Department I for Internal Medicine,
 84 University Hospital of Cologne, Cologne/Germany and German
 85 Centre for Infection Research, partner site Bonn-Cologne,
 86 Germany, ²⁸A.O.U. Policlinico Vittorio Emanuele Catania,
 87 Biometec – University of Catania/Italy, ²⁹Infectious Diseases
 88 Unit, S. Orsola-Malpighi, Department of Medical and Surgical
 89 Sciences, University of Bologna, Bologna/Italy, ³⁰Department
 90 of Clinical Microbiology and Immunology, Sackler School of
 91 Medicine, Tel Aviv University, Tel Aviv/Israel, ³¹Institute of
 92 Medical Microbiology, University Hospital Essen, University of
 93 Duisburg- Essen, Essen/Germany, ³²Department of of
 94 Microbiology, Faculty of Medicine and Dentistry, Palacky
 95 University Olomouc and University Hospital Olomouc/Czech

96 Republic, ³³Clinical Microbiology, Puerta del Mar University
 97 Hospital, University of Cádiz, Cádiz/Spain, ³⁴Infectious Diseases
 98 Unit, 3rd Department of Pediatrics, Faculty of Medicine,
 99 Aristotle University School of Health Sciences, Hippokration
 100 General Hospital, Thessaloniki/Greece, ³⁵Department of
 101 Medical Microbiology, Hacettepe University Medical School,
 102 Ankara/Turkey, ³⁶Division of Mycology, Department of Medial
 103 Microbiology, Chandigarh/India, ³⁷Escola Paulista de Medicina,
 104 Federal University of São Paulo, São Paulo/Brazil,
 105 ³⁸Departamento de Micología, Instituto de Medicina Regional,
 106 Universidad Nacional del Nordeste, CONICET,
 107 Resistencia/Argentina, ³⁹ Division of Clinical Mycology,
 108 Department of Microbiology, Vall d'Hebron University
 109 Hospital, Barcelona, ⁴⁰Department of Medical Mycology and
 110 Parasitology/Invasive Fungi Research Center, Mazandaran
 111 University of Medical Sciences, Sari/Iran, ⁴¹School of Medicine,
 112 European University Cyprus, Nicosia/Cyprus, ⁴²Department of
 113 Clinical Mycology, Allergy and Immunology, North Western
 114 State Medical University, Saint Petersburg/Russia,
 115 ⁴³Department I for Internal Medicine, University Hospital of
 116 Cologne, Cologne/Germany, ⁴⁴CBS-KNAW Fungal Biodiversity
 117 Centre, Utrecht/The Netherlands, ⁴⁵Hacettepe University
 118 Medical School, Department of Infectious Diseases and Clinical
 119 Microbiology, Ankara/Turkey, ⁴⁶Department of Medical

120 Statistics, Informatics and Health Economics, Medical
121 University of Innsbruck, Innsbruck/Austria, ⁴⁷Department of
122 Dermatology, Hospital Ernest Lluch Martin, Calatayud,
123 Zaragoza/Spain
124
125 Running title: TerrNet Study
126
127 Corresponding author: Cornelia Lass-Flörl, Division of Hygiene
128 and Medical Microbiology. University of Innsbruck,
129 Schöpfstraße 41, 6020 Innsbruck, Austria
130 Tel.No. 0043 512 9003-70703, Fax No. 0043 512 9003-73700
131 email: cornelia.lass-floerl@i-med.ac.at
132
133 Length of abstract : 87 words
134 Length of paper (excluding abstract and ref.) : 1172 words

135 **Abstract**

136 Objectives: A prospective international multicentre

137 surveillance study was conducted to investigate the

138 prevalence and amphotericin B (AMB) susceptibility of

139 *Aspergillus terreus* species complex infections.

140 Methods: Three hundred seventy cases from 21 countries

141 were evaluated.

142 Results: The overall prevalence of *A. terreus* species complex

143 among patients investigated and with mold positive cultures

144 was 5.2% (370/7116). AMB MICs were ranging from 0.125 to

145 32 mg/L, (median 8mg/L).

146 Conclusions: *A. terreus* species complex infections cause a

147 wide spectrum of aspergillosis and the majority of cryptic

148 species display high AMB MICs.

149 Introduction:

150 *Aspergillus terreus* species complex holds an exceptional
151 position within the aspergilli, as it appears to be a rare
152 pathogen of infection and displays polyene resistance [1,2,3].
153 *A. terreus* is a common cause of invasive aspergillosis (IA) at
154 the M. D. Anderson Cancer Center in Houston, USA, and the
155 University Hospital of Innsbruck, Austria [3,4,5]. Almost no
156 data are available on how frequently this species occurs
157 elsewhere and whether differences within amphotericin B
158 (AMB) susceptibility exist. Our objective was to investigate the
159 global prevalence of *A. terreus* species complex in fungal
160 diseases and to survey AMB susceptibility.

161

162 Methods:

163 An international surveillance network was established on
164 behalf of the European Fungal Infection Study Group, the
165 International Society for Human and Animal Mycology
166 *Aspergillus terreus* working group, and the European
167 Confederation of Medical Mycology. 38 centres from 21
168 countries participated. Each centre collected isolates and
169 reported the number of *A. terreus* and fungal pathogens
170 detected for 12 consecutive months (2014-2015). Patient
171 characteristics', epidemiological data, and antifungal
172 treatment were documented through an online questionnaire

173 using www.clinicalsurveys.net online platform. Patients were
174 classified according to the European Organisation for the
175 Research and Treatment of Cancer/Mycoses Study Group
176 consensus definitions [6] by the participating centres. Unless
177 otherwise noted, the isolation of *A. terreus* from sputa of non-
178 neutropenic patients was categorized as colonisation. Isolates
179 were sent to the Division of Hygiene and Medical Microbiology
180 for molecular species identification [7,8] and susceptibility
181 testing according to EUCAST (European Committee on
182 Antimicrobial Susceptibility Testing) method [2]. *A. terreus*
183 strains were identified to the cryptic species level by
184 sequencing partial beta-tubulin and applying a validated in-
185 house database owned by Jos Houbraken, CBS Fungal
186 Biodiversity Center, Utrecht, The Netherlands. An AMB
187 epidemiological cut-off value of 4mg/L was set for *A. terreus*
188 [2].

189 This study was approved by the Ethics Commission of the
190 Medical University of Innsbruck (UN4926).

191 Results:

192 461 cases were enrolled of which 91 were excluded because of
193 insufficient patient documentation (n=45) or lack of fungal
194 isolates (n=46) being available. Consequently, this survey
195 comprises 370 eligible cases with an equal number of
196 corresponding *A. terreus* isolates. Cases derived from Europe

197 (n=261), followed by Middle East (n=70), India (n=19), South
 198 America (n=10), and North America (n=10) (Figure 1). *A.*
 199 *terreus* sensu stricto (n=315), *A. citrinoterreus* (n=36), *A.*
 200 *alabamensis* (n=6), *A. hortai* (n=10), *A. floccosus* (n=1), and *A.*
 201 *neoafricanus* (n=1) were identified. One isolate (*A. terreus*
 202 1214) was most close to *A. alabamensis* and might represent a
 203 new species. Thus, cryptic species accounted for 14.9%
 204 (55/370) with *A. citrinoterreus* (36/55, 65.5%) being dominant.
 205 AMB MICs ranged from 0.125 to 32 mg/L for *A. terreus* sensu
 206 stricto; MICs for all cryptic species were consistently higher,
 207 ranging from 2 to 32 mg/L, see Table 1. According to the
 208 EUCAST cut-off values, 194 isolates (52.4%) were classified as
 209 non-wild types. A proportion of 6.3% (n=20) of the *A. terreus*
 210 sensu stricto isolates displayed lower MICs, ranging from 0.25
 211 – 0.5 mg/L. Isolates were predominantly acquired from Spain
 212 (n=85) and Austria (n=49), see Figure 1.
 213 Underlying diseases e described in Table 2. Species distribution
 214 did not differ per underlying disease and specimen
 215 investigated (Table 2). Diseases comprised IA (25.1%), allergic
 216 broncho-pulmonary aspergillosis (12.4%), chronic aspergillosis
 217 (11.4%), COPD exacerbation (5.5%), aspergilloma (3.7%), otitis
 218 externa (2.5%), and wound infections (0.7%). 25.1% and 27.3%
 219 of the patients suffered from proven and probable IA, 28.6%
 220 were colonized, 10.1% had onychomycosis, and 8.9% had

221 mycological documented diseases such as otitis externa,
222 aspergilloma and others.
223 Using a random effects model the pooled estimated
224 proportion was 5.6% (95% CI 3.8 to 7.7) with $I^2 = 92\%$
225 ($p < 0.0001$) and the proportions ranged from 0.0% to 58.3%.
226 These calculations were done with MedCalc 16.8.4. Four
227 reference centres and one centre dealing with onychomycosis
228 only were excluded from the analysis.
229 A total of 68 patients received antifungal treatment at the
230 time of fungal diagnosis, 12 were treated with AMB or
231 liposomal - AMB. The remaining 56 received combinations of
232 azoles and echinocandins and improved. Only one patient died
233 due to the *A. terreus* infection. No information on outcome
234 was available in 13 patients.

235

236 Discussion:

237 Infections due to *A. terreus* species complex were detected in
238 21 countries and 38 centres with an overall prevalence of 5.2%
239 among mold infections. High AMB MICs were frequently
240 observed and crossed all cryptic species. Infections were
241 reported from all over the world with three main specific
242 findings. Firstly, Spain and Austria were the countries with the
243 highest density of *A. terreus* isolates collected. Secondly, the
244 number of *A. terreus* cases enrolled varied from centre to

245 centre, and displayed a broad range from zero to several cases
246 per country. Thirdly, it seems that few susceptible AMB
247 variants exist within *A. terreus* sensu stricto.
248 Taking into account the differences on the environmental
249 conditions, host related characteristics, and the use of
250 antifungal agents, it is not possible to conclude on the
251 particular biogeography of *A. terreus* species complex. In
252 addition, one has to be aware that data collected may depend
253 on the quality of care, patient demographics, infection control
254 practices, frequency of specimen collection, and laboratory
255 methodology. Hence, further studies are needed to determine
256 whether specific risk and/or environmental factors are
257 associated with infections by *A. terreus*.
258 Notable was the fact that *Aspergillus* section *Terrei* was most
259 commonly isolated from patients suffering from chronic lung
260 diseases (39.2%). No similar data have been reported [10] and
261 it remains to be seen whether *A. terreus* reflects an emerging
262 pathogen of this disease entity.
263 *A. terreus* is a poor target for AMB and hence is reported as
264 resistant [2]. The role of isolates with MICs <0.5 mg/L needs
265 further evaluation. The pharmacodynamic target may be
266 attained with the standard AMB dose for isolates with MICs
267 ≤ 0.25 mg/l [10] and infections were successfully treated with
268 high dose liposomal-AMB [11].

269 Cryptic species accounted for 14.8%, with *A. citrinoterreus*
270 being the most prevalent. Although the clinical implications of
271 sibling species of *A. terreus* are less well understood, our study
272 confirms that these species are generally resistant to AMB and
273 are causing a wide spectrum of invasive and non-invasive
274 aspergillosis. Guinea et al. [12] observed *A. citrinoterreus*
275 acting mainly as a co-pathogen with *A. fumigatus*.
276 Our study has some limitations. We do not have a
277 comprehensive worldwide *A. terreus* survey network and
278 some countries are missing for a variety of reasons. Also,
279 generally, the diagnosis of fungal infections is difficult to
280 obtain and may often be based on detection of biomarkers
281 rather than on isolation of the infecting organism. Hence,
282 some cases may have been missed and chronic lung diseases
283 were not specified in more detail. Further, we have no data
284 available on co-infections which may complicate diseases. The
285 centres included represent a convenience sample. However,
286 this is the largest and geographically most diverse study on the
287 contemporary epidemiology of *A. terreus* species complex
288 infections worldwide.

289 Our study shows that *A. terreus* sensu stricto is widely
290 distributed in climatically divergent countries, and that cryptic
291 species display high AMB MICs. *A. terreus* species complex was

292 most commonly isolated from patients suffering from chronic
293 lung diseases (39.2%).

ACCEPTED MANUSCRIPT

294 Funding

295 This work was supported by ECMM, ISHAM and EFISG and in
296 part by an unrestricted research grant through the Investigator
297 Initiated Studies Program of Astellas and Pfizer as well as by
298 the Austrian Science Fund (W1253-B24 doctoral program
299 HOROS).

300

301

302 Transparency Declaration

303 We declare that we have no conflicts of interest related to this
304 study.

305 CLF received research grants, travel grants or honorarium as a
306 speaker or advisor from Gilead Sciences, Pfizer, Schering

307 Plough, MSD and Basilea, VAA received research grants or

308 honorarium as a speaker or advisor from Astellas, Pfizer and

309 Schering Plough, MSD. WB received honorarium as a speaker

310 from Pfizer, Schering Plough and MSD. CF received research

311 grants or honorarium as a speaker or advisor from Astellas,

312 Gilead Sciences, MSD and Basilea. AH received travel grants

313 from Astellas and served on the advisory board of Gilead. EJ

314 received travel grants or honorarium as a speaker or advisor

315 from Bio-Rad, Gilead Sciences, Pfizer, Schering Plough, and

316 MSD. DPK has received research support from Pfizer, Astellas

317 Pharma, honoraria from Astellas, Merck, Pfizer and he serves

318 on the advisory boards of Merck, Amplyx, Cidara and F2G. JFM
319 received grants from Astellas, Basilea and Merck. He has been
320 a consultant to Astellas, Basilea and Merck and received
321 speaker's fees from Merck, United Medical and Gilead. ALC
322 received educational grants from Pfizer, Gilead and United
323 Medical and research grant from Pfizer, Astellas. MCA received
324 research grants/contract work (Paid to SSI) from Amplyx,
325 Basilea, Cidara, F2G, Gilead, Pfizer & T2Candida, speaker
326 honoraria from Astellas, Basilea, Gilead, MSD, Novartis, Pfizer
327 & T2Candida. NK has received research support from Pfizer,
328 honoraria from Astellas, Merck and Pfizer, and he serves on
329 the advisory boards of Merck and Pfizer. OAC is supported by
330 the German Federal Ministry of Research and Education, has
331 received research grants from Actelion, Aranis, Astellas,
332 AstraZeneca, Basilea, Bayer, Cidara, Duke University (NIH
333 UM1AI104681), F2G, Gilead, GSK, Leeds University, MedPace,
334 Melinta Therapeutics, Merck/MSD, Miltenyi, Pfizer, Rempex,
335 Roche, Sanofi Pasteur, Scynexis, Seres Therapeutics, The
336 Medicine Company, is a consultant to Anacor, Amplyx,
337 Actelion, Astellas, Basilea, Cidara, Da Volterra, F2G, Gilead,
338 Inositec AG, Janssen Pharmaceuticals, Matinas, Menarini,
339 Merck/MSD, Paratek Pharmaceuticals, Scynexis, Seres,
340 Summit, Vical, Vifor, and received lecture honoraria from
341 Astellas, Basilea, Gilead and Merck/MSD. BW received

342 research support from Pfizer, travel grants or honorarium as a
343 speaker or advisor from Gilead Sciences, Pfizer, Astellas, MSD
344 and Basilea. LK grant from Gilead and has been an adviser to
345 Astellas, Gilead, Schering-Plough, has received research grants
346 from Gilead, Merck, Sharpe&Dohme, Schering-Plough and has
347 received honoraria for educational lectures from Gilead,
348 Pfizer, Merck, Sharpe & Dohme, Schering-Plough and Janssen.
349 AMT received research support from Gilead Sciences, travel
350 grants or honorarium as speaker from Gilead Sciences, Pfizer,
351 Astellas. KL has received research grants from Gilead, MSD and
352 Pfizer, received travel support from MSD, Pfizer and Gilead
353 and received lecture honoraria from Gilead, MSD, and Pfizer.
354 SAA received Investigator Initiated Research grant support
355 from Pfizer and speaker honoraria or travel grants from
356 Astellas, Gilead, Merck and Pfizer. SO received research grants
357 from Pfizer and Astellas, honorarium as advisor from MSD,
358 Astellas Pharma and Gilead Sciences. ER reports grants, and
359 non-financial support from Astellas, Pfizer, Gilead, and Merck,
360 outside the submitted work. AAI received research and/or
361 travel grants from Gilead Sciences, F2G and Pfizer. OU was
362 consultant for and has received travel grants from Merck,
363 Astellas, Pfizer, and Gilead. S.M.H. has received research and
364 travel grants from Astellas, Gilead, and MSD, research grants
365 from Basilea, and travel grants from Pfizer; was consultant to

- 366 Basilea; and received honoraria as a speaker from Astellas and
367 Merck.

ACCEPTED MANUSCRIPT

368 **References**

- 369 1. Perkhofer S, Lass-Flörl C, Hell M, Russ G, Krause R, Hönigl
370 M, et al. The Nationwide Austrian *Aspergillus* Registry: a
371 prospective data collection on epidemiology, therapy and
372 outcome of invasive mould infections in
373 immunocompromised and/or immunosuppressed patients.
374 Int J Antimicrob Agents 2010;36:531-6.
- 375 2. Arendrup MC, Cuenca-Estrella M, Lass-Flörl C, Hope WW;
376 European Committee on Antimicrobial Susceptibility
377 Testing Subcommittee on Antifungal Susceptibility Testing
378 (EUCAST-AFST). EUCAST technical note on *Aspergillus* and
379 amphotericin B, itraconazole, and posaconazole. Clin
380 Microbiol Infect. 2012;18:E248-50.
- 381 3. Lass-Flörl C, Grif K, Mayr A, Petzer A, Gastl G, Bonatti H et
382 al. Epidemiology and outcome of infections due to
383 *Aspergillus terreus*: 10-year single center experience. Br J
384 Haematol. 2005;131:201-207.
- 385 4. Lass-Flörl C, Grif K, Kontoyiannis DP. Molecular typing of *A.*
386 *terreus* isolates collected in Houston, Texas and Innsbruck,
387 Austria: evidence of great genetic diversity. J Clin
388 Microbiol. 2007;45:2686-90.
- 389 5. Lackner M, Coassin S, Haun M, Binder U, Kronenberg F,
390 Haas H et al. Geographically predominant genotypes of
391 *Aspergillus terreus* species complex in Austria: a

- 392 microsatellite typing study. Clin Microbiol Infect.
393 2016;22:270-276.
- 394 6. De Pauw B, Walsh TJ, Donnelly JP, Stevens DA, Edwards JE,
395 Calandra T et al. Revised definitions of invasive fungal
396 disease from the European Organization for Research and
397 Treatment of Cancer/Invasive Fungal Infections
398 Cooperative Group and the National Institute of Allergy
399 and Infectious Diseases Mycoses Study Group
400 (EORTC/MSG) Consensus Group. Clin Infect Dis
401 2008;46:1813-21.
- 402 7. Balajee S A, Baddley JW, Peterson SW, Nickle D, Varga J,
403 Boey A et al., ISHAM Working Group on *A. terreus*.
404 *Aspergillus alabamensis*, a new clinically relevant species in
405 the section *Terrei*. Eukaryotic Cell 2009;8:713–22.
- 406 8. Kathuria S, Sharma C, Singh PK, Agarwal P, Agarwal K,
407 Hagen F et al. Molecular epidemiology and in-vitro
408 antifungal susceptibility of *Aspergillus terreus* species
409 complex isolates in Delhi, India: Evidence of genetic
410 diversity by amplified fragment length polymorphism and
411 microsatellite typing. PLoS ONE 2015;10:e0118997.
- 412 9. Denning DW, Cadranel J, Beigelman-Aubry C, Ader F,
413 Chakrabarti A, Blot S et al. Chronic pulmonary aspergillosis:

- 414 rationale and clinical guidelines for diagnosis and
415 management. Eur Respir J. 2016;47:45-68.
- 416 10. Elefanti A, Mouton JW, Verweij, PE, Zerva L, Meletiadis J.
417 Susceptibility breakpoints for amphotericin B and
418 *Aspergillus* species in an in vitro pharmacokinetic-
419 pharmacodynamic model simulating free-drug
420 concentrations in human serum. Antimicrob Agents
421 Chemother. 2014;58:2356-2362.
- 422 11. Lewis RE, Albert NP, Liao G, Wang W, Prince RA,
423 Kontoyiannis DP. High-dose induction liposomal
424 amphotericin B followed by de-escalation is effective in
425 experimental *Aspergillus terreus* pneumonia. J Antimicrob
426 Chemother. 2013;68:1148-1151.
- 427 12. Guinea J, Sandoval-Denis M, Escibano P, Peláez T, Guarro
428 J, Bouza E. *Aspergillus citrinoterreus*, a new species of
429 section Terrei isolated from samples of patients with non-
430 hematological predisposing conditions. J Clin Microbiol.
431 2015;53:611-617.

Table 1. Distribution of amphotericin B MICs against *Aspergillus terreus* species complex isolates collected during the study period and tested according to EUCAST methodology

Species	Amphotericin B MICs, mg/L								
	0.125	0.25	0.5	1	2	4	8	16	32
<i>A. terreus</i> sensu stricto	3	7	10	14	36	81	86	55	23
<i>A. citrinoterreus</i>					3	13	8	7	5
<i>A. hortai</i>					1	2	5	2	
<i>A. alabamensis</i>					2	3	1		
<i>A. floccosus</i>						1			
<i>A. neoaffricanus</i>									1
Potential new species							1		

Table 2. Species distribution of *Aspergillus terreus* species complex isolated from the various human specimens

Species	Specimens, total numbers						
	Sputa	Bronchoalveolar lavages and tracheal secretions	Body-fluids	Biopsies	Swabs	Others	Total
<i>A. terreus</i> sensu stricto	126	65	53	33	17	21	315
<i>A. citrinoterreus</i>	14	7	3	5	3	4	36
<i>A. hortai</i>	4	2			1	3	10
<i>A. alabamensis</i>	3	2			1		6
<i>A. floccosus</i>					1		1
<i>A. neoaffricanus</i>						1	1
Potential new species				1			1
Total	147	76	56	39	23	29	370


* aspirates, wound secretions, nails

Fig. 1a - c. Overview of countries and *Aspergillus terreus* species complex isolated numbers collected during the study period:


a) *Aspergillus citrinoterreus*


b) *Aspergillus terreus* sensu stricto

c) *Aspergillus hortai*

Aspergillus citrinoterreus

Total=36

Aspergillus terreus sensu stricto

Aspergillus hortai

Total=10